

Emerging nanotechnologies for cancer immunotherapy

Sourabh Shukla^{1,2} and Nicole F Steinmetz^{1,2,3,4,5}

¹Department of Biomedical Engineering, Case Western Reserve University, Cleveland, OH 44106, USA; ²Case Comprehensive Cancer Center, Case Western Reserve University, Cleveland, OH 44106, USA; ³Department of Radiology, Case Western Reserve University, Cleveland, OH 44106, USA; ⁴Department of Materials Science and Engineering, Case Western Reserve University, Cleveland, OH 44106, USA; ⁵Department of Macromolecular Science and Engineering, Case Western Reserve University, Cleveland, OH 44106

Corresponding authors: Sourabh Shukla. Email: sourabh.shukla@case.edu; Nicole F Steinmetz. Email: nicole.steinmetz@case.edu

Abstract

Founded on the growing insight into the complex cancer-immune system interactions, adjuvant immunotherapies are rapidly emerging and being adapted for the treatment of various human malignancies. Immune checkpoint inhibitors, for example, have already shown clinical success. Nevertheless, many approaches are not optimized, require frequent administration, are associated with systemic toxicities and only show modest efficacy as monotherapies. Nanotechnology can potentially enhance the efficacy of such immunotherapies by improving the delivery, retention and release of immunostimulatory agents and biologicals in targeted cell populations and tissues. This review presents the current status and emerging trends in such nanotechnology-based cancer immunotherapies including the role of nanoparticles as carriers of immunomodulators, nanoparticles-based cancer vaccines, and depots for sustained immunostimulation. Also highlighted are key translational challenges and opportunities in this rapidly growing field.

Keywords: Cancer immunotherapy, nanotechnology, vaccines, immunomodulation

Experimental Biology and Medicine 2016; 241: 1116–1126. DOI: 10.1177/1535370216647123

Introduction

Cancer immunotherapies – revitalizing the immune system against cancer

One of the major developments in the fight against cancer is the emergence of immunotherapies that are aimed at harnessing the exquisite and specific power of the immune system against malignancies.^{1,2} The renewed excitement and push for novel immunotherapies stems from the success of two different strategies – adaptive T-cell therapy based on chimeric antigen receptors (CARs) and checkpoint blockade. The former, CAR T cells therapy, still under clinical evaluation, is based on genetically engineering patient's own T-cells with CARs that recognizes tumor antigens. These CAR T cells are then expanded *in vitro* and infused back in patients, where they are likely to recognize and kill cancer cells.³ Checkpoint blockade therapies, on the other hand work by inhibiting pathways that keep the duration and strength of immune system in check.⁴ The recent approval of two checkpoint blockade therapies targeting the receptors CTLA-4 and PD-1 have come on the back of several successful clinical trials where treatment with checkpoint blockade inhibitors has resulted in striking T cell function restoration in melanoma, renal cell carcinoma and lung cancer.^{4,5}

These developments and other similar efforts are clearly fueled by growing insights into the nature and consequences of interactions between tumors and the immune system, which frequently impede the development and function of anti-tumor immune response.^{6–8} The challenge for immunotherapies is to modulate such interactions towards successful recognition and elimination of cancer cells. The durability and specificity of such strategies has the potential to generate long-lived therapeutic effects with limited systemic toxicities.^{9,10} However, growing evidences suggest that immunotherapies could be most beneficial as an adjuvant therapy to conventional chemo- and radiation therapies.¹¹

The premise of an immune system-mediated intervention in cancer progression lies in the capacity of the immune system to distinguish between self and non-self.^{12,13} While highly equipped and effective in the eradication of pathogens, the ability of the immune system to effectively deal with transforming cancer cells is hampered by the fact that the cancer cell's origin is self, and because of depletion of self-antigen reactive T lymphocytes during development.¹⁴ Nevertheless, cancer is characterized by the accumulation of a variable number of genetic alterations and the loss of normal cellular regulatory processes, resulting in expression of neo-antigens arising from mutated

genes, chromosomal aberrations or overexpression of embryonic antigens.^{15,16} These neo-antigens differentiate cancer from normal cells and enable their immune-recognition, as indicated by the presence of basal levels of tumor antigen-specific cellular and humoral responses in subsets of cancer patients.^{13,17} However, a highly immunosuppressive microenvironment under the influence of cytokines such as TGF- β and IL-2,¹⁸ along with immunosuppressive cells such as Foxp3⁺ T_{regs} (regulatory T cells),¹⁹ myeloid-derived suppressor cells (MDSCs)²⁰ and M2-type macrophages,²¹ or combinations thereof, keeps such autologous immune response in check. Moreover, through intricate immunoediting mechanisms, including antigen shedding, negative selection of antigenic cancer cells, down-regulation of MHC-I molecules and turning off activated T cells via negative regulators such as PD-1,⁴ cancer cells evade immunosurveillance and the tumor prevails.⁷

To overcome these hurdles and tip the scale in favor of anti-tumor immune response, a diverse set of immunotherapeutic approaches have been explored. While adoptive T cell transfer (ACT)²² and cancer vaccines²³ are aimed at boosting tumor reactive immune cell populations, cytokines, immunomodulatory antibodies and small molecule drugs have been employed to overcome the immunosuppressive tumor microenvironment.²⁴ The goal of cancer immunotherapy is to initiate or reinitiate a self-sustaining cycle of cancer immunity, enabling it to amplify and propagate.² Although numerous such strategies have been explored, only a handful of them have been approved and adapted for clinical use with only a modest rate of success.¹ Thus, there is a huge scope for further development in terms of specificity, enhanced effectiveness and reduced toxicities.

Many immunotherapeutic strategies currently in pre-clinical or clinical evaluation are based on traditional drug development approaches where nanomedicine has already made significant contributions by improving stability, bio-distribution through targeted delivery, bioavailability and efficacy of cytotoxic drugs or imaging contrast agents.^{25,26} Ongoing clinical evaluations of several candidate nanoformulations in conjugation with a wide range of chemotherapeutic or immunotherapeutic payloads are a testament of the efficacy of nanoparticle-based drug delivery approaches.^{27,28} For example, Nanoplatin NC-6004, a cisplatin containing polymeric nanocarrier is under Phase I/II clinical trial,²⁹ whereas cyclodextrin-based nanoparticles CALAA-01 that delivers small-interfering RNA (siRNA) agent to shut down a key enzyme (ribonucleotide reductase) in cancer cells is under Phase I evaluation.³⁰ Similarly, an engineered adenovirus nanoparticle-based drug delivery platform is under Phase-I dose escalation study for delivery of cancer immunotherapy to patients with chronic lymphocytic leukemia (CLL)³¹ while the pH sensitive polymeric nanoparticle CRLX101 loaded with camptothecin is undergoing Phase II clinical trials.³²

Based on those very possibilities, an excellent opportunity for nanotechnology-mediated refinement exists in the field of immunotherapy.^{33–35} This review details some of the key immunotherapeutic strategies and highlights nanotechnology-based interventions that are being pursued to improve

the overall efficacy of such approaches, as summarized in Figure 1. Additionally, critical barriers to the successful translation of these emerging technologies are also discussed.

Nanocarriers to deliver tumor microenvironment immunomodulators

A number of immunomodulatory and immunostimulatory molecules such as cytokines, chemokines and targeted antibodies have been identified for their important roles in countering the highly immunosuppressive tumor microenvironment. Cytokine IL-2 promotes proliferation of effector functions of cytotoxic T lymphocytes (CTLs) and has shown clinical efficacy in malignant melanoma and renal carcinoma.³⁶ IL-2 has also resulted in enhanced efficiency of other immunotherapies.³⁷ Other cytokines such as IL-21 and IL-18 modulate both innate and adaptive immune responses through activation of CD4⁺/CD8⁺ T cells, natural killer (NK) cells, and B cells while suppressing T_{reg} cells.^{38,39} Systemic administration of IL-21 and IL-18 also leads to enhanced production of IFN- γ , IL-2, tumor necrosis factor- α (TNF- α), granulocyte macrophage colony-stimulating factor (GM-CSF), IL-1 β and IL-6 by activating T cells. Similarly, type I interferons (IFN- α and β) demonstrate anti-tumor activities through stimulation of NK cell activity and suppression of allospecific suppressor T cells. Indeed, the administration of type I interferons has shown promise and efficacy in clinical trials in the setting of leukemia, melanoma and renal cell carcinoma.⁴⁰ Type II interferons (IFN- γ) induce apoptosis, upregulate HLA-I and HLA-II and therefore promote antigen presentation in cancer cells, which in turn mediates tumor rejection and has shown efficacy in the setting of ACT therapies.⁴¹ Other non-specific immunomodulators such as Toll-like receptor (TLR7/9) agonists (e.g., synthetic oligonucleotides CpG) promote Th1 polarization, trigger activation of innate and adaptive immune responses, lead to dendritic cell (DC) activation and proliferation of CD4⁺/CD8⁺ T cells and modulate suppressive functions of T_{reg} cells.^{42,43}

Even though the immunomodulatory effects of these small molecules in overturning the suppressive tumor microenvironment are well documented, drawbacks are associated with such therapies: Besides the short half-life, stability and bioavailability challenges akin to many conventional therapeutic candidates, systemic toxicities of cytokines arising due to their broad spectrum of biological activity on a wide variety of cells are major safety concerns.^{44–46} Cytokines could lead to non-specific lymphocyte activation in circulation and increased incidences of autoimmune and allergic responses. IL-1, IL-2, IL-6, TNF, and TGF- β could lead to modulation of hepatic metabolisms.⁹ Similarly, systemic administration of CD-40 agonist used to trigger CD40 signaling for activation of antigen presenting cells can lead to widespread symptoms of cytokine release syndrome, ocular inflammation, elevated levels of hepatic enzymes, and hematologic toxicities including T-cell depletion. Besides, agonistic anti-CD40 therapy has also been linked to long-term immunosuppression mediated by activation-induced apoptosis of CD4⁺ and CD8⁺ T cells.^{47,48} Similarly, overexposure to CpG could result in suppression

of adaptive T cell immunity.^{49,50} Likewise, IL-2 administration at high doses causes vascular leak syndrome (VLS; also known as capillary leak syndrome), which is associated with increased vascular permeability, hypotension, pulmonary edema, liver cell damage and renal failure.⁵¹

Other side effects of IL-2 are hypothyroidism, thrombocytopenia, anemia, coagulopathy, or impairment of neutrophil chemotaxis, autoimmunity, neurotoxicity and myocarditis.^{52,53} In addition, the ability of IL-2 to stimulate T_{Reg} cells diminishes the beneficial effects of stimulating tumor-specific T cell response.⁵⁴ Systemic administration of IL-2 with adaptively transferred T cells could also cause multi organ failures in severe cases. At the same time, dose dependent toxicities including thrombocytopenia, fatigue, and pyrexia have been associated with checkpoint blockade inhibitors.⁹

To overcome these challenges and provide a pathway for safe clinical use of immunomodulatory cytokines/chemokines, nanotechnology approaches hold great promise for the path forward. Nanoformulations consisting of such immunomodulatory molecules have improved bioavailability due to significantly prolonged circulation times of the carrier particles and *in vivo* stability of payload against serum inactivation and enzyme degradation.^{55,56} For example, intravenous administrations of liposomes containing cytokines such as IFN- γ , IFN- α , IL-2 or TNF- α enhance the plasma residence time.^{55,57,58} On the other hand, intraperitoneal, intramuscular, subcutaneous or intranasal administration of cytokine carrying liposomes and polymeric particles creates local depots and increases residence times of the immunostimulatory payloads at the site (discussed in details later).^{59–61} Additionally, requirement of external or physiological stimuli ensures the release of immunostimulatory cargo only at targeted sites and further improve its bioavailability and safety.^{62–65}

Specifically, nanoparticle-based delivery promotes the preferential accumulation and retention of immunomodulators in tumors due to the enhanced permeability and retention (EPR) effect, while minimizing off-target systemic toxicities, thereby improving potential for clinical translation of such therapies.^{66,67} Building on EPR effect-mediated nanoparticle homing, nanotechnologies are undergoing development targeting and modulating the immunosuppressive tumor microenvironment to attain efficacy of immunotherapies. For example, based on the passive tumor homing properties, lipid-coated calcium phosphate nanoparticles (LCP-NPs) have been used for tumor microenvironment immunomodulation by delivering TGF- β siRNA and thereby down-regulating the levels immunosuppressive TGF- β within the tumor. LCP-NPs have also been used to deliver a broad spectrum anti-inflammatory triterpenoid – methyl-2-cyano-3,12-dioxooleana-1,9(11)-dien-28-oate (CDDO-Me) – that significantly reduced T_{Reg} and MDSC populations. The delivery of immunostimulatory molecules by LCP-NP has been combined with vaccination strategies using an LCP vaccine delivering a tumor antigen (Trp 2 peptide) and adjuvant (CpG oligonucleotide) to DCs – the combination therapy resulted in improved efficacy over vaccine only treatments.^{68,69}

Similarly, EPR-mediated accumulation of liposome-encapsulated polymer nanogels has been utilized for intratumoral delivery of cytokine (IL-2) and TGF- β receptor I inhibitor – SB505124, a hydrophobic small molecule drug, leading to inhibition of TGF- β receptor I and subsequent expansion of T cells and NK cells by blocking key immunosuppressive pathways.⁷⁰ Similarly, by delivering PD-L1 siRNA using polyethylenimine (PEI) liposomes, PD-L1 levels have been knocked down leading to immunosuppressive to immunostimulatory phenotype changes in human and mouse ovarian cancer-associated DCs with subsequent increase in tumor-reactive CD8⁺ T cell numbers and improved mice survival.⁷¹ Liposomal delivery of IL-2 has also showed enhanced therapeutic effects with reduced toxicities in a variety of other tumors including liver and lung cancers leading to significant reduction in tumor growth.^{72,73}

In addition to their use for systemic delivery and homing to tumors, nanoparticle formulations show benefits for intratumoral delivery of immunomodulators: based on their size, the nanoparticle-based formulation of immunotherapies limits their escape into systemic circulation, thus minimizing off-target effects and maximizing local immunostimulation. For example, immunostimulatory liposomes conjugated with IL-2 and anti-CD137 antibodies targeting activated T cells led to increased IL-2 dosing within the tumor when delivered directly via intratumoral vs. systemic injections. The intratumoral treatment resulted in a higher ratio of tumor-infiltrating CD8⁺ T cells over regulatory T cells in established melanomas.⁴⁶ Likewise, PEGylated liposome formulation have been used to deliver agonistic anti-CD40 antibodies and TLR agonist CpG molecules using intratumoral administration resulting in significant tumor inhibition while sequestering the immunostimulatory payload in targeted tissues and reducing its systemic leakage, thus minimizing off-target inflammatory effects.⁴⁵ Similarly, intratumoral administration of CpG payloads on gold nanoparticles has shown to induce significant macrophage and DC infiltration in tumors and significantly affected tumor growth by concentrating the CpG oligonucleotides in the tumor tissue and lowering the high dose requirements of systemic administrations.⁷⁴

Passive tumor homing and intratumoral administration of nanoparticles are accompanied by their natural tropism towards phagocytic cells of the innate immune system *in vivo* including monocytes, neutrophils, macrophages and dendritic cells. Such affinity towards immune cells can also be used to deliver immunomodulating payloads to tumor-infiltrating immune cell populations and thereby reprogramming the tumor microenvironment. For example, nanocomplexes encapsulating CpG oligonucleotide and anti-IL-10 and anti-IL-10 receptor antisense oligonucleotides were efficiently captured by tumor-associated macrophages (TAMs) and resulted in altered macrophage phenotypes, leading to a significant anti-tumor effect in a hepatoma murine model.⁷⁵ To improve the partition in TAMs over macrophages associated with the mononuclear phagocyte system (MPS), mannose-modified polymeric micelles containing acid-sensitive PEG modifications were developed. The PEG shielding reduced uptake by the MPS

in circulation at neutral pH and improved tumor accumulation through persistent circulation. The acidic pH in the tumor resulted in PEG shedding and uptake by TAMs and their subsequent reprogramming.⁷⁶

However, nanoparticles have also been targeted to immune cells in circulation to deliver payloads to the tumor tissue. RGD-targeted single walled carbon nanotubes (SWCNTs) have showed enhanced tumor accumulation via hitchhiking Ly6C^{hi} monocytes in the circulation that are recruited to the site of the tumor in response to inflammation.⁷⁷ Also, in a recent study, gold nanoparticles combining mouse vascular endothelial growth factor (VEGF)-siRNA with TAM-targeting M2 peptide have been used to inhibit both TAMs and cancer cells by targeting the VEGF pathway in both cell populations. Such synergistic inhibitory effects on both cancer cells and the immunosuppressive TAM population resulted in significant tumor regression and disease control for extended periods in orthotopic lung cancer in mice.⁷⁸ These examples highlight the utility of nanoparticles to target phagocytic cells for immune activation; tissue- and cell-specificity can be further tailored by altering nanoparticle shape, size, charge, hydrophobicity and surface chemistry.^{33,79,80}

In addition to their natural interactions with mononuclear phagocyte cells, nanoparticles carrying immunostimulatory payloads have also been utilized to modulate the functioning of T and B cells for therapeutic interventions. Adaptive T cell immunotherapy plays a central role in cancer therapy; cancer antigen-specific T cells can be expanded using vaccines (*in vivo*) or through ACT, the latter requires the *ex vivo* expansion of T cells followed by infusion into patient. While effective, ACT has many drawbacks because the methods are expensive, cumbersome and personalized.²² To enhance the technology, in one study, dextran-coated iron oxide particles with surface-coupled MHC-Ig dimers and anti-CD28 antibodies were designed to allow magnetic field-based aggregation of particles bound to T cell receptors (TCRs). *Ex vivo* stimulation of T cells with these particles in the presence of a magnetic field-enhanced TCR clustering reduced the threshold of activation of T cells and improved the efficacy of adaptive T cell therapy.⁸¹ Moreover, nanotechnology opens the door for the *in vivo* targeting, priming and expansion of T cells. For example, *in vivo* loading of T cells with lipid nanoparticle “backpacks” carrying stimulatory cytokines was demonstrated. The nanoparticle-mediated *in vivo* priming resulted in 80-fold increased T cell expansion and significant enhancements in the efficacy of ACT without systemic toxicity.⁸² Similarly, circulating adaptive T cells were targeted *in vivo* by IL-2 loaded liposomes via anti-Thy1 antibodies, resulting in enhanced T cell proliferation more effectively compared to administration of soluble cytokines.⁸³ These approaches overcome a decline in function of transplanted T cells following infusion, particularly in the setting of solid cancers with a highly immunosuppressive microenvironment.

Finally, we recently demonstrated the use of virus-based nanoparticles as an immunotherapeutic, where the properties of the nanoparticle itself unlocked a potent anti-tumor immune response. We have demonstrated that plant-

derived virus-like particles stimulate a potent immune-mediated anti-tumor response when introduced into the tumor microenvironment after tumors are established: VLPs from cowpea mosaic virus (CPMV), without nucleic acids, LPS, or any other recognized immune adjuvants, generated an effective anti-tumor immune response in mouse models of multiple tumor types, including triple negative breast cancer, disseminated ovarian cancer and primary and metastatic melanoma. The particles are not cytolytic to tumor cells and the effects are immune mediated. Most importantly, preliminary data indicate that the effect is systemic and durable, resulting in immune memory protecting mice from re-challenge.⁸⁴ The immunotherapy follows an *in situ* vaccination approach in which immune-stimulatory reagents (here CPMV) are applied directly into the suspected metastatic site or into an identified primary tumor. This approach modulates the local microenvironment to relieve immunosuppression and potentiate anti-tumor immunity against antigens expressed by the tumor. This approach not only offers the potential for new therapeutics but also may lead to new levels of understanding how the immune system defines “danger signals”.

Improving cancer vaccines – mediators of adaptive immune response

A wide range of cancer vaccines has been evaluated for a variety of human malignancies.^{85,86} The overarching goal is to deliver tumor-associated antigens to professional antigen presenting cells (APCs) to elicit adaptive immune responses mediated by tumor-specific cytotoxic T cells and antibodies. As an active immunotherapeutic approach aimed at stimulating endogenous anti-tumor immune response, cancer vaccines offer effective long-term protection against recurring and residual tumors. However, development of an effective therapeutic vaccine against established disease is challenging, and despite decades of pursuit, establishment of successful vaccination strategies based on proteins, peptides, autologous dendritic cells or tumor cells have largely been unsuccessful.^{18,87} While vaccines based on autologous cells are costly and technically challenging, peptide-based cancer vaccination suffers from inefficient uptake, processing and presentation of the delivered epitopes by activated professional APCs.^{23,51,87–90} Moreover, such vaccination strategies have led to the generation of low avidity tumor-specific T cell responses. Whole protein vaccination with powerful and often poorly tolerated adjuvants, immunostimulatory cytokines such as IL-2 or GM-CSF and/or TLR agonists have failed to induce clinically significant anti-tumor responses.^{91–93} However, spontaneous tumor antigen-specific and high avidity T cell response has been shown to control tumor progression. Together with the demonstration of significantly reduced tumor burden in both patients and animal models following ACT, this suggests that high avidity tumor-specific CTL response could lead to long-lasting immunoprotection against relapsing or residual cancer.^{85,86,94}

Nanoparticle-based vaccine approaches offer multiple advantages that could fulfill the stringent requirements for the generation of such high avidity tumor-reactive

T cells.^{95,96} Presenting antigens/epitopes on nanoparticulate carriers not only provides the requisite stability and longevity, it also facilitates efficient interactions with key immune cell populations.^{88,97} Clearly, nanoparticle engineering principles for vaccine platforms are different from those targeted for delivery of immunomodulators. While the former seeks interaction with APCs and other phagocytes, the latter aims to generally avoid phagocytic clearance thus prolonging circulation and improved tumor penetration. Nanoparticle-based vaccine formulations can improve the resulting immunostimulation by promoting multivalent receptor cross-linking, by altering intracellular processing and presentation or by colocalizing synergistic cues from the antigen, adjuvants and costimulatory molecules within the same cellular populations. The particulate nature of nanoparticles mimic pathogen-associated molecular patterns (PAMPs) that are perceived as danger signals and drive protective immunity.⁹⁸ Such patterns are recognized by the pattern recognizing receptors (PRRs) such as TLRs on immune cells, specifically APCs, and facilitate enhanced uptake of nanoparticle-based vaccines by these cells.^{99,100} Activation of PRRs provides immunogenic cues to the immune system instructing it to launch specific response to the antigens carried by the nanoformulation.

B cells have evolved to recognize multivalent display of antigens on microbial surfaces and play a major role in vaccine-mediated antibody responses, thus enhancement of their engagement is crucial for immunotherapies based on cancer vaccines. Nanoparticle-based antigen display that mimic pathogen structural features are highly efficient in engaging B cell receptors (BCRs) to promote greater signaling, antigen internalization and processing of antigens for presentation to CD4⁺ T cells.¹⁰¹ Multivalent display of antigens on nanoparticles also leads to TLR stimulation promoting strong humoral responses with long-lived high avidity antibody responses mediated by secretion of co-stimulatory cytokines such as IFN- α and IL-12.¹⁰² For example, we have demonstrated that potato virus X (PVX) displaying HER2-derived B-cell epitopes effectively generates HER2-specific antibodies.¹⁰³ Similarly, plant viral nanoparticles coupled to a weak idiotype (Id) tumor antigen have been used as a conjugate vaccine to induce antibody formation against a murine B-cell malignancy.¹⁰⁴ Other virus-like particles have similarly been evaluated as efficient vaccine carriers.^{105–108} For a detailed review, we would like to refer the reader to our recent article (Lee, 2016).¹⁴⁷

Professional APCs, particularly dendritic cells, are critical initiators of adaptive immune response, comprising both humoral and cellular responses, and are therefore an important target for anti-cancer nanomedicine. Nanoparticle-based vaccines are readily taken up by DCs and are associated with enhanced anti-tumor response as compared to soluble antigens.¹⁰⁹ Following processing in endolysosomal compartments, soluble exogenous antigens are exclusively presented on MHC-II molecules to activate CD4⁺ helper T cells that in turn stimulate B cell-mediated antibody response. However, via a process called cross-presentation, nanoparticle-mediated delivery of antigenic peptides also results in cytosolic release of antigens where

MHC-I loading can occur, resulting in CD8⁺ T cell priming and ensuing cytotoxic T cell response.¹¹⁰ Such cross-presentation of cancer antigens have been demonstrated for a wide range of distinct nanoparticles and new strategies are being developed to improve the efficiency of such cross-presentation.^{111–113} These include strategies employed for cytosolic drug delivery of membrane-impermeable molecules such as via endolysosomal disruption through the proton sponge effect using biodegradable nanogels,¹¹⁴ endolysosomolytic and pH-responsive micelles,¹¹⁵ as well as endoplasmic reticulum (ER) targeting approaches where nanoparticles shuttle to cytosol following endosome-ER fusion.¹¹⁶

A significant advantage of nanoformulations is the control over their transport kinetics that facilitates tissue-specific delivery of antigens. Tumor antigens drain into tumor draining lymph nodes (TDLNs), where they are taken up by professional APCs, including DCs leading to subsequent presentation and priming of T cells. TDLNs are rich in phenotypically and functionally immature APCs and are therefore key targets for priming APCs using cancer vaccines and the subsequent adaptive immune response. The size-dependent lymphatic drainage of nanoformulations has been established and is an important design consideration for developing cancer vaccines: Nanoparticles between 20 and 45 nm optimally drain to lymph nodes and are retained.¹¹⁷ While smaller particles are likely to be flushed out of lymph nodes, those larger than 100 nm are less efficiently transported from the peripheral injection sites, generally via cell-mediated transport.¹¹⁸ Upon subcutaneous injection, only a small fraction of the vaccine is delivered to DCs whereas the majority is cleared by the body or engulfed by other immune cells.

By targeting DCs, vaccine efficiencies can be further improved by overcoming non-specific uptake of nanoparticle-based vaccines. For example, cancer vaccines based on biodegradable poly(lactic-co-glycolic acid) (PLGA) nanoparticles when coated with an agonistic α CD40-mAb (NP-CD40), showed highly efficient and selective delivery to DCs *in vivo* and improved priming of CD8⁺ T cells against two independent tumor-associated antigens.¹¹⁹ Additionally, targeting a specific subset of DCs could also define the type of stimulated immune response.¹²⁰ For example, TLR7 and TLR9 agonists can convert the tolerogenic plasmacytoid DCs to innate immunostimulatory types, whereas targeting various C-type lectins can modulate variable adaptive response. For instance, DC-SIGN, DEC-205, DNNGR-1 and Langerin favors CD8⁺ T cell cellular (Th1) responses while CD4⁺ and B cell humoral (Th2) responses are achieved by targeting of DCIR2.¹²¹ The targeting of multiple DCs subtypes, in particular, holds great potential to enhance vaccine efficiencies.

Sustained immunostimulation through *in situ* depots and artificial APCs

Nanoparticles-based depots have recently gained attraction as a source of sustainable immune stimulus. For example, cytokine depots composed of liposomes or polymeric

particles carrying pro-inflammatory cytokines have been developed for anticancer vaccines and for intratumoral administration for therapy. These cytokine-loaded particles can enhance tumor-specific immune response in conjugation with tumor antigen from irradiated tumor cells or co-encapsulated with the cytokines.⁵⁵ Using these strategies, GM-CSF encapsulated in polymer particles or IFN- γ /IL-2 in liposomes caused increased leukocyte infiltration/increased humoral response and enhanced cytolytic capacity of CD8⁺ T cells resulting in higher fraction of mice surviving melanoma challenge.¹²² Similar studies extended in human trials involved incorporation of tumor-specific idotype isolated from follicular lymphoma patient into IL-2 containing liposomes and monthly vaccinations which resulted in sustained tumor-specific CD4⁺ and CD8⁺ T cell response and continuous remission.¹²³

On the other hand, cytokine depots have been also employed to treat primary tumors with peri- or intratumoral injections. Here, primary tumors serve as the source of antigen while cytokine depot activate leukocytes in tumor microenvironment and promote immunotherapy against primary tumor and metastasized tumor cells. Local injection of polymeric microparticles loaded with IL-2 for the treatment of brain or liver tumors has showed that this approach was more effective at treating tumors and protecting against rechallenge than tumor cells engineered to express IL-2.¹²⁴ Compared to treatment with soluble IL-2, liposomal IL-2 treatment of B16 melanoma bearing mice resulted in higher survival rates, slower tumor growth rates prior to resection and increased recruitment and protected mice against rechallenge with melanoma.¹²⁴

Further studies using this strategy have involved particles with combinations of various cytokines including IL-12, TNF- α , GM-CSF or IL-18. Treatment of established tumors with IL-12 loaded particles prior to surgical resections promoted systemic anti-tumor immunity that prevented recurrence and metastasis.¹²⁵ Furthermore, particles with combination of IL-12 and GM-CSF showed eradication of primary tumors through CD4⁺ and CD8⁺ cells, while the effect on metastasis was through NK/NKT cells.¹²⁶

The concept of immunotherapeutic depot has also been similarly used in therapeutic vaccine DepoVaxTM (DPX-0907), which employs a liposome-based platform harboring custom formulated mixtures of CD8⁺ T-cell peptide epitopes, a tetanus toxoid derived Th epitope, and an adjuvant of choice (such as a toll-like receptor agonist) to provide signals for improved antigen presentation. The liposomes carry incorporated hydrophilic antigens and adjuvant directly into an oil medium such as Montanide ISA51 VG, entrapping all vaccine ingredients in a form suitable for efficient uptake, processing and presentation by antigen-presenting cells (APCs). Such DPX formulated vaccines have been shown to induce effective immune responses after a single-dose administration.¹²⁷

Another development in this area is the design and engineering of artificial APCs (aAPCs) that are synthetic mimics of natural antigen presenting cells, to promote T cell activation and subsequent expansion, both *ex vivo* and *in vivo*. Essentially, aAPCs are particles to which proteins

required for T cell activation, such as MHC-epitope complexes, agonist anti-CD3 and agonist anti-CD28, have been conjugated.¹²⁸ Both spatial and temporal organization of these signals during aAPC/T cell contact is important for efficient T cell activation. The first generation aAPCs were composed of solid, micron-sized polystyrene beads or with iron oxide cores and were used for *ex vivo* expansion of T cells. Their large size provided a large area of contact between aAPCs and T cells. However, the second generation of aAPCs engineered for *in vivo* applications were smaller particles at the nanometer size scale (<100 nm) that showed favorable distribution to T-cell-rich regions such as spleens and lymph nodes upon systemic administration.¹²⁹ Shape is again a key design parameter here. CD8⁺ T cells migrated preferentially to the long axis of ellipsoidal aAPCs and the extended length of contact resulted in enhanced proliferation and anti-tumor response.¹³⁰

Challenges and opportunities for nanotechnology-based immunotherapies

Despite almost a decade-long research, preclinical and clinical, into the immunostimulatory potential of nanotechnology-based platforms, only a handful of approaches thus far have received clinical approval. A success story in this regard is T-VEC. An attenuated version of herpes simplex virus, T-VEC is a genetically engineered oncolytic virus that specifically replicates in cancer cells and leads to anti-cancer response by secreting cytokine GM-CSF. With successful phase III trials, T-VEC has been recently approved for treatment of melanoma patients as an injectable formulation.¹³¹

However, the translational gap highlights not only the complex relationship of cancer and immunology, but also underlines poor understanding of the *in vivo* behavior of nanoparticles and its safety concerns. Undesirable immunotoxicity of nanoparticles, adverse interactions with biomolecules, long-term accumulation at off target sites, remain challenging. Generation of pro-inflammatory cytokines (such as IL-6 and TNF- α) and inflammasome response contribute to immunotoxicity. For example, mesoporous hollow silica nanoparticles and carbon nanotubes have been associated with the induction of pro-inflammatory cytokines, liver damage and activation of Kupffer cells.^{132,133} Similarly, nanosize TiO₂ particles lead to oxidative stress, neutrophil activation and inflammation in lungs,¹³⁴ while carbon nanotubes have been implicated for inflammatory fibrogenic pulmonary response.¹³⁵ On the other hand, MPS clearance resulting from adsorption of serum proteins, complements and immunoglobulins results in off-target accumulations.¹³⁶

As highlighted in the above discussions, the combination of physical and chemical properties of nanomaterials influences their biological interactions and hence their applicability for specific immunotherapeutic interventions. Therefore, depending on the immunostimulatory pathways and mechanisms to be perturbed, the nanoparticle platform and route of administration must be selected based on overall biodistribution, pharmacokinetics, range of cellular interactions and toxicological risk assessments. Extensive

preclinical evaluation of various nanoparticle platforms, both *in vitro* and *in vivo*, is therefore critical.

Some key aspects of the physiological behavior of nanoparticles that have been driving forces behind their applications in nanomedicine are still fairly ineffective.¹³⁷ Delivering therapeutic payload to tumors via systemic administration remains challenging. The EPR effect is critical for nanoparticle homing into solid tumors for drug delivery or immunomodulation of the tumor microenvironment.⁶⁷ However, adequacy of the EPR effect remains a controversial subject amongst the scientific community and varies greatly between various cancer types and from primary tumors to metastatic sites. EPR is also a function of nanocarrier morphology and results in only a small fraction homing into tumors; a large fraction is still non-specifically delivered to off-target sites and could lead to severe side effects.¹³⁸ Even after reaching tumor sites, penetration of nanoparticle therapeutics in tumor mass may be impaired because of barriers created by abnormal tumor physiology

including abnormal tumor structures, such as physically compromised vasculature, abnormal ECM, and high interstitial fluid pressure.

Targeting tumor vasculatures or cancer cell receptors could increase uptake and retention of nanoparticles in endothelial and cancer cells. However, from an immunotherapeutic point of view, targeting immunomodulators to immune cell populations within the tumor stroma could be a more effective strategy as even if a small population of these cells is activated, they would proliferate through a cascade of specific mechanisms leading to effective coordinated adaptive anti-tumor response. Also, keeping in mind the broad spectrum of immune cells that could potentially be activated by therapeutic immunomodulators, systemic toxicity remains a significant concern. Concentrating the immunotherapeutic drugs into nanoparticles followed by tissue- and cell-targeted delivery has been shown to overcome dose-limiting toxicities by minimizing off-target accumulation. Similarly, improving nanoparticle stability in




Figure 1 Cancer immunotherapeutic strategies and nanotechnology-based intervention to overcome challenges of current immunotherapy approaches. (A color version of this figure is available in the online journal.)

circulation, upon serum protein conjugations and under varying pH and redox potentials should also be key to rule out premature release of such broad-spectrum cargo.¹³⁹ At the same time, emphasizing the development of safer biodegradable nanomaterials will facilitate earlier translation of such platform technologies.

As briefly mentioned earlier, design principles for nanoparticle-based vaccines targeting cancer antigen-specific adaptive immune responses are essentially contrasting to those for delivery of therapeutic immunomodulators. The former requires effective uptake and processing by professional antigen presenting cells in secondary lymphoid tissues, while the latter specifically requires avoidance of phagocytosis by APCs. Thus, selection of platform technology becomes critical. Also, given the consistently evolving nature of the tumor, multi-epitope vaccines are likely to be more effective because combinations of antigens may reduce the window for the tumor to down-regulate the antigens and escape immune system detection.¹⁴⁰ Therefore, nanoparticles that facilitate presentation of high payloads of multiple epitopes are preferable. Cancer vaccines are also likely to be more efficient under conditions of minimal residual disease or in combination with immunomodulators reducing the immune suppression of established tumors.

Nanomanufacturing and quality control and assurance must be considered along the translational pathway. Improved scaled-up production of nanoparticles of interest with excellent reproducibility and consistency of composition, physical properties and chemical addressability is a critical bottleneck for a large range of contemporary nanoparticle platform technologies. Advances such as PRINTTM technology¹⁴¹ and biology-inspired materials such as plant virus-based technologies that can be manufactured in plants¹⁴² are some of the examples where such criteria have been met. Protein-based platform technologies based on plant viruses and bacteriophages have added benefits of genetic programmability, whereby genes for proteins, targeting peptides and antigenic motifs of interests can be inserted into the viral genome and expressed as a fusion product to the coat proteins.^{143,144} Similarly, specific chemical functionalities in the form of amino acid side chains can be coded into the viral genome with precise spatial control to expand the contemporary virus-based and virus-like nanoparticle libraries with altered reactivities.¹⁴³ Once genetically engineered in such manner, large-scale production of identical clones through propagation in their natural plant and bacterial hosts provides excellent control over the nanomanufacturing process, leading to excellent structural and functional monodispersity. Moreover, engineering interventions during the self-assembly process of such viral-based materials also provides another level of control over their structural and functional properties by altering their aspect ratios/size¹⁴⁵ or by loading non-natural cargos within the viral capsids.¹⁴⁶ Such control over the structure-function relationship is currently unattainable with most synthetic nanomaterials and is immensely desirable towards development of improved platform technologies.

In summary, cancer immunotherapy holds great promise in cancer therapy. Collaboration between cancer immunologists and engineers will further the understanding of the

complex and underlying immunology, therefore driving technological development. With emerging nanotechnological interventions, efficacy of many such immunotherapies could be drastically improved and a merger of these two rapidly growing fields of science could facilitate clinical translation of many cancer immunotherapies.

Authors' contributions: SS and NFS both contributed to planning the review outline, collecting literature and writing the mini-review.

ACKNOWLEDGEMENTS

This work was supported by a grant from the Susan G. Komen foundation (CCR14298962 to NFS).

DECLARATION OF CONFLICTING INTERESTS

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

REFERENCES

- Pardoll DM. Immunology beats cancer: a blueprint for successful translation. *Nat Immunol* 2012;**13**:1129–32
- Mellman I, Coukos G, Dranoff G. Cancer immunotherapy comes of age. *Nature* 2011;**480**:480–9
- Grupp SA, Kalos M, Barrett D, Aplenc R, Porter DL, Rheingold SR, Teachey DT, Chew A, Hauck B, Wright JF, Milone MC, Levine BL, June CH. Chimeric antigen receptor-modified T cells for acute lymphoid leukemia. *N Engl J Med* 2013;**368**:1509–18
- Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer* 2012;**12**:252–64
- Brower V. Checkpoint blockade immunotherapy for cancer comes of age. *J Natl Cancer Inst* 2015; **107**:4–6
- Gajewski TF, Schreiber H, Fu YX. Innate and adaptive immune cells in the tumor microenvironment. *Nat Immunol* 2013;**14**:1014–22
- Schreiber RD, Old LJ, Smyth MJ. Cancer immunoediting: integrating immunity's roles in cancer suppression and promotion. *Science* 2011;**331**:1565–70
- Lindau D, Gielen P, Kroesen M, Wesseling P, Adema GJ. The immunosuppressive tumour network: myeloid-derived suppressor cells, regulatory T cells and natural killer T cells. *Immunology* 2013;**138**:105–15
- Weber JS, Yang JC, Atkins MB, Disis ML. Toxicities of immunotherapy for the practitioner. *J Clin Oncol* 2015;**33**:2092–9
- Alatrash G, Jakher H, Stafford PD, Mittendorf EA. Cancer immunotherapies, their safety and toxicity. *Expert Opin Drug Saf* 2013;**12**:631–45
- Lake RA, Robinson BW. Immunotherapy and chemotherapy – a practical partnership. *Nat Rev Cancer* 2005;**5**:397–405
- Jiang H, Chess L. How the immune system achieves self-nonself discrimination during adaptive immunity. *Adv Immunol* 2009;**102**:95–133
- Swann JB, Smyth MJ. Immune surveillance of tumors. *J Clin Invest* 2007;**117**:1137–46
- Zou W. Regulatory T cells, tumour immunity and immunotherapy. *Nat Rev Immunol* 2006;**6**:295–307
- Linley AJ, Ahmad M, Rees RC. Tumour-associated antigens: considerations for their use in tumour immunotherapy. *Int J Hematol* 2011;**93**:263–73
- Bright RK, Bright JD, Byrne JA. Overexpressed oncogenic tumor-self antigens. *Hum Vaccin Immunother* 2014;**10**:3297–3305
- Vesely MD, Kershaw MH, Schreiber RD, Smyth MJ. Natural innate and adaptive immunity to cancer. *Annu Rev Immunol* 2011;**29**:235–71

18. Sheng J, Chen W, Zhu HJ. The immune suppressive function of transforming growth factor-beta (TGF-beta) in human diseases. *Growth Factors* 2015;**33**:92-101
19. Oleinika K, Nibbs RJ, Graham GJ, Fraser AR. Suppression, subversion and escape: the role of regulatory T cells in cancer progression. *Clin Exp Immunol* 2013;**171**:36-45
20. Gabrilovich DI, Nagaraj S. Myeloid-derived suppressor cells as regulators of the immune system. *Nat Rev Immunol* 2009;**9**:162-74
21. Chanmee T, Ontong P, Konno K, Itano N. Tumor-associated macrophages as major players in the tumor microenvironment. *Cancers (Basel)* 2014;**6**:1670-90
22. Rosenberg SA, Restifo NP. Adoptive cell transfer as personalized immunotherapy for human cancer. *Science* 2015;**348**:62-68
23. Melero I, Gaudernack G, Gerritsen W, Huber C, Parmiani G, Scholl S, Thatcher N, Wagstaff J, Zielinski C, Faulkner I, Mellstedt H. Therapeutic vaccines for cancer: an overview of clinical trials. *Nat Rev Clin Oncol* 2014;**11**:509-24
24. Mahoney KM, Rennert PD, Freeman GJ. Combination cancer immunotherapy and new immunomodulatory targets. *Nat Rev Drug Discov* 2015;**14**:561-84
25. Mitragotri S, Anderson DG, Chen X, Chow EK, Ho D, Kabanov AV, Karp JM, Kataoka K, Mirkin CA, Petrosko SH, Shi J, Stevens MM, Sun S, Teoh S, Venkatraman SS, Xia Y, Wang S, Gu Z, Xu C. Accelerating the translation of nanomaterials in biomedicine. *ACS Nano* 2015;**9**:6644-54
26. Peer D, Karp JM, Hong S, Farokhzad OC, Margalit R, Langer R. Nanocarriers as an emerging platform for cancer therapy. *Nat Nanotechnol* 2007;**2**:751-60
27. Bourzac K. Nanotechnology: Carrying drugs. *Nature* 2012;**491**:S58-60
28. Ventola CL. The nanomedicine revolution: part 2: current and future clinical applications. *P T* 2012;**37**:582-91
29. Plummer R, Wilson RH, Calvert H, Boddy AV, Griffin M, Sludden J, Tilby MJ, Eatock M, Pearson DG, Ottley CJ, Matsumura Y, Kataoka K, Nishiya T. A Phase I clinical study of cisplatin-incorporated polymeric micelles (NC-6004) in patients with solid tumours. *Br J Cancer* 2011;**104**:593-8
30. Davis ME, Zuckerman JE, Choi CH, Seligson D, Tolcher A, Alabi CA, Yen Y, Heidel JD, Ribas A. Evidence of RNAi in humans from systemically administered siRNA via targeted nanoparticles. *Nature* 2010;**464**:1067-70
31. Castro JE, Melo-Cardenas J, Urquiza M, Barajas-Gamboa JS, Pakbaz RS, Kipps TJ. Gene immunotherapy of chronic lymphocytic leukemia: a phase I study of intranodally injected adenovirus expressing a chimeric CD154 molecule. *Cancer Res* 2012;**72**:2937-48
32. Zuckerman JE, Gritti I, Tolcher A, Heidel JD, Lim D, Morgan R, Chmielowski B, Ribas A, Davis ME, Yen Y. Correlating animal and human phase Ia/Ib clinical data with CALAA-01, a targeted, polymer-based nanoparticle containing siRNA. *Proc Natl Acad Sci USA* 2014;**111**:11449-54
33. Smith DM, Simon JK, Baker JR Jr. Applications of nanotechnology for immunology. *Nat Rev Immunol* 2013;**13**:592-605
34. Sheng WY, Huang L. Cancer immunotherapy and nanomedicine. *Pharm Res* 2011;**28**:200-14
35. Goldberg MS. Immunoengineering: how nanotechnology can enhance cancer immunotherapy. *Cell* 2015;**161**:201-4
36. Nicholas C, Lesinski GB. Immunomodulatory cytokines as therapeutic agents for melanoma. *Immunotherapy* 2011;**3**:673-90
37. Rosenberg SA. IL-2: the first effective immunotherapy for human cancer. *J Immunol* 2014;**192**:5451-8
38. Davis MR, Zhu Z, Hansen DM, Bai Q, Fang Y. The role of IL-21 in immunity and cancer. *Cancer Lett* 2015;**358**:107-114
39. Tian H, Shi G, Yang G, Zhang J, Li Y, Du T, Wang J, Xu F, Cheng L, Zhang X, Dai L, Chen X, Zhang S, Yang Y, Yu D, Wei Y, Deng H. Cellular immunotherapy using irradiated lung cancer cell vaccine co-expressing GM-CSF and IL-18 can induce significant antitumor effects. *BMC Cancer* 2014;**14**:48
40. Zitvogel L, Galluzzi L, Kepp O, Smyth MJ, Kroemer G. Type I interferons in anticancer immunity. *Nat Rev Immunol* 2015;**15**:405-14
41. Tagawa M, Kawamura K, Li Q, Tada Y, Hiroshima K, Shimada H. A possible anticancer agent, type III interferon, activates cell death pathways and produces antitumor effects. *Clin Dev Immunol* 2011;**2011**:479013
42. Shiota H, Klinman DM. Use of CpG oligonucleotides for cancer immunotherapy and their effect on immunity in the tumor microenvironment. *Immunotherapy* 2013;**5**:787-9
43. Klinman DM. Immunotherapeutic uses of CpG oligodeoxynucleotides. *Nat Rev Immunol* 2004;**4**:249-58
44. Vanneman M, Dranoff G. Combining immunotherapy and targeted therapies in cancer treatment. *Nat Rev Cancer* 2012;**12**:237-51
45. Kwong B, Liu H, Irvine DJ. Induction of potent anti-tumor responses while eliminating systemic side effects via liposome-anchored combinatorial immunotherapy. *Biomaterials* 2011;**32**:5134-47
46. Kwong B, Gai SA, Elkhader J, Wittrup KD, Irvine DJ. Localized immunotherapy via liposome-anchored Anti-CD137 + IL-2 prevents lethal toxicity and elicits local and systemic antitumor immunity. *Cancer Res* 2013;**73**:1547-58
47. Bartholdy C, Kauffmann SO, Christensen JP, Thomsen AR. Agonistic anti-CD40 antibody profoundly suppresses the immune response to infection with lymphocytic choriomeningitis virus. *J Immunol* 2007;**178**:1662-70
48. Berner V, Liu H, Zhou Q, Alderson KL, Sun K, Weiss JM, Back TC, Longo DL, Blazar BR, Wiltout RH, Dai L, Chen X, Zhang S, Yang Y, Yu D, Wei Y, Deng H. IFN-gamma mediates CD4+ T-cell loss and impairs secondary antitumor responses after successful initial immunotherapy. *Nat Med* 2007;**13**:354-60
49. Heikenwalder M, Polymenidou M, Junt T, Sigurdson C, Wagner H, Akira S, Zinkernagel R, Aguzzi A. Lymphoid follicle destruction and immunosuppression after repeated CpG oligodeoxynucleotide administration. *Nat Med* 2004;**10**:187-192
50. Wingender G, Garbi N, Schumak B, Jungerkes F, Endl E, von Bubnoff D, Steitz J, Striegler J, Moldenhauer G, Tuting T, Heit A, Huster KM, Takikawa O, Akira S, Busch DH, Wagner H, Hämmerling GJ, Knolle PA, Limmer A. Systemic application of CpG-rich DNA suppresses adaptive T cell immunity via induction of IDO. *Eur J Immunol* 2006;**36**:12-20
51. Donohue JH, Rosenberg SA. The fate of interleukin-2 after in vivo administration. *J Immunol* 1983;**130**:2203-8
52. Atkins MB, Mier JW, Parkinson DR, Gould JA, Berkman EM, Kaplan MM. Hypothyroidism after treatment with interleukin-2 and lymphokine-activated killer cells. *N Engl J Med* 1988;**318**:1557-63
53. Klempner MS, Noring R, Mier JW, Atkins MB. An acquired chemotactic defect in neutrophils from patients receiving interleukin-2 immunotherapy. *N Engl J Med* 1990;**322**:959-65
54. Boyman O, Surh CD, Sprent J. Potential use of IL-2/anti-IL-2 antibody immune complexes for the treatment of cancer and autoimmune disease. *Expert Opin Biol Ther* 2006;**6**:1323-31
55. Christian DA, Hunter CA. Particle-mediated delivery of cytokines for immunotherapy. *Immunotherapy* 2012;**4**:425-41
56. Petros RA, DeSimone JM. Strategies in the design of nanoparticles for therapeutic applications. *Nat Rev Drug Discov* 2010;**9**:615-27
57. van der Veen AH, Eggermont AM, Seynhaeve AL, van T, ten Hagen TL. Biodistribution and tumor localization of stealth liposomal tumor necrosis factor-alpha in soft tissue sarcoma bearing rats. *Int J Cancer* 1998;**77**:901-6
58. Kedar E, Gur H, Babai I, Samira S, Even-Chen S, Barenholz Y. Delivery of cytokines by liposomes: hematopoietic and immunomodulatory activity of interleukin-2 encapsulated in conventional liposomes and in long-circulating liposomes. *J Immunother* 2000;**23**:131-45
59. Kedar E, Palgi O, Golod G, Babai I, Barenholz Y. Delivery of cytokines by liposomes. III. Liposome-encapsulated GM-CSF and TNF-alpha show improved pharmacokinetics and biological activity and reduced toxicity in mice. *J Immunother* 1997;**20**:180-193
60. Epstein DA, Stewart WE 2nd. Altered pharmacological properties of liposome-associated human interferon-alpha. *J Virol* 1982;**41**:575-82
61. Anderson PM, Katsanis E, Sencer SF, Hasz D, Ochoa AC, Bostrom B. Depot characteristics and biodistribution of interleukin-2 liposomes: importance of route of administration. *J Immunother* (1991) 1992;**12**:19-31

62. Yuyama Y, Tsujimoto M, Fujimoto Y, Oku N. Potential usage of thermosensitive liposomes for site-specific delivery of cytokines. *Cancer Lett* 2000;**155**:71–7
63. ten Hagen TL, Seynhaeve AL, van Tiel ST, Ruiter DJ, Eggermont AM. Pegylated liposomal tumor necrosis factor- α results in reduced toxicity and synergistic antitumor activity after systemic administration in combination with liposomal doxorubicin (Doxil) in soft tissue sarcoma-bearing rats. *Int J Cancer* 2002;**97**:115–120
64. Heffernan MJ, Kasturi SP, Yang SC, Pulendran B, Murthy N. The stimulation of CD8⁺ T cells by dendritic cells pulsed with polyketal microparticles containing ion-paired protein antigen and poly(inosinic acid)-poly(cytidylic acid). *Biomaterials* 2009;**30**:910–18
65. Haining WN, Anderson DG, Little SR, von Bergwelt-Baildon MS, Cardoso AA, Alves P, Kosmatopoulos K, Nadler LM, Langer R, Kohane DS. pH-triggered microparticles for peptide vaccination. *J Immunol* 2004;**173**:2578–85
66. Davis ME, Chen ZG, Shin DM. Nanoparticle therapeutics: an emerging treatment modality for cancer. *Nat Rev Drug Discov* 2008;**7**:771–82
67. Maeda H, Nakamura H, Fang J. The EPR effect for macromolecular drug delivery to solid tumors: Improvement of tumor uptake, lowering of systemic toxicity, and distinct tumor imaging in vivo. *Adv Drug Deliv Rev* 2013;**65**:71–79
68. Xu Z, Wang Y, Zhang L, Huang L. Nanoparticle-delivered transforming growth factor- β siRNA enhances vaccination against advanced melanoma by modifying tumor microenvironment. *ACS Nano* 2014;**8**:3636–45
69. Zhao Y, Huo M, Xu Z, Wang Y, Huang L. Nanoparticle delivery of CDDO-Me remodels the tumor microenvironment and enhances vaccine therapy for melanoma. *Biomaterials* 2015;**68**:54–66
70. Park J, Wrzesinski SH, Stern E, Look M, Criscione J, Ragheb R, Jay SM, Demento SL, Agawu A, Licona Limon P, Ferrandino AF, Gonzalez D, Habermann A, Flavell RA, Fahmy TM. Combination delivery of TGF- β inhibitor and IL-2 by nanoscale liposomal polymeric gels enhances tumour immunotherapy. *Nat Mater* 2012;**11**:895–905
71. Cubillos-Ruiz JR, Engle X, Scarlett UK, Martinez D, Barber A, Elgueta R, Wang L, Nesbeth Y, Durant Y, Gewirtz AT, Sentman CL, Kedl R, Conejo-Garcia JR. Polyethylenimine-based siRNA nanocomplexes reprogram tumor-associated dendritic cells via TLR5 to elicit therapeutic antitumor immunity. *J Clin Invest* 2009;**119**:2231–44
72. Kanaoka E, Takahashi K, Yoshikawa T, Jizomoto H, Nishihara Y, Uchida N, Maekawa R, Hirano K. A significant enhancement of therapeutic effect against hepatic metastases of M5076 in mice by a liposomal interleukin-2 (mixture). *J Control Release* 2002;**82**:183–7
73. Neville ME, Robb RJ, Popescu MC. In situ vaccination against a non-immunogenic tumour using intratumoural injections of liposomal interleukin 2. *Cytokine* 2001;**16**:239–50
74. Lin AY, Almeida JP, Bear A, Liu N, Luo L, Foster AE, Drezek RA. Gold nanoparticle delivery of modified CpG stimulates macrophages and inhibits tumor growth for enhanced immunotherapy. *PLoS One* 2013;**8**:e63550
75. Huang Z, Zhang Z, Jiang Y, Zhang D, Chen J, Dong L, Zhang J. Targeted delivery of oligonucleotides into tumor-associated macrophages for cancer immunotherapy. *J Control Release* 2012;**158**:286–92
76. Zhu S, Niu M, O'Mary H, Cui Z. Targeting of tumor-associated macrophages made possible by PEG-sheddable, mannose-modified nanoparticles. *Mol Pharm* 2013;**10**:3525–30
77. Smith BR, Ghosn EE, Rallapalli H, Prescher JA, Larson T, Herzenberg LA, Gambhir SS. Selective uptake of single-walled carbon nanotubes by circulating monocytes for enhanced tumour delivery. *Nat Nanotechnol* 2014;**9**:481–7
78. Conde J, Bao CC, Tan YQ, Cui DX, Edelman ER, Azevedo HS, Byrne HJ, Artzi N, Tian FR. Dual targeted immunotherapy via in vivo delivery of biohybrid RNAi-peptide nanoparticles to tumor-associated macrophages and cancer cells. *Adv Funct Mater* 2015;**25**:4183–94
79. Champion JA, Mitragotri S. Shape induced inhibition of phagocytosis of polymer particles. *Pharm Res* 2009;**26**:244–9
80. Kumar S, Anselmo AC, Banerjee A, Zakrewsky M, Mitragotri S. Shape and size-dependent immune response to antigen-carrying nanoparticles. *J Control Release* 2015;**220**(Pt A): 141–8
81. Perica K, Tu A, Richter A, Bieler JG, Edidin M, Schneck JP. Magnetic field-induced T cell receptor clustering by nanoparticles enhances T cell activation and stimulates antitumor activity. *ACS Nano* 2014;**8**:2252–60
82. Stephan MT, Moon JJ, Um SH, Bershteyn A, Irvine DJ. Therapeutic cell engineering with surface-conjugated synthetic nanoparticles. *Nat Med* 2010;**16**:1035–41
83. Zheng Y, Stephan MT, Gai SA, Abraham W, Shearer A, Irvine DJ. In vivo targeting of adoptively transferred T-cells with antibody- and cytokine-conjugated liposomes. *J Control Release* 2013;**172**:426–35
84. Lizotte PH, Wen AM, Sheen MR, Fields J, Rojasopondist P, Steinmetz NF, Fiering S. In situ vaccination with cowpea mosaic virus nanoparticles suppresses metastatic cancer. *Nat Nanotechnol* 2015
85. Irvine DJ, Hanson MC, Rakhra K, Tokatlian T. Synthetic nanoparticles for vaccines and immunotherapy. *Chem Rev* 2015;**115**:11109–46
86. Bolhassani A, Safaiyan S, Rafati S. Improvement of different vaccine delivery systems for cancer therapy. *Mol Cancer* 2011;**10**:3
87. Azvolinsky A. Cancer vaccines: always a bridesmaid, never a bride? *J Natl Cancer Inst* 2013;**105**:248–9
88. Krishnamachari Y, Geary SM, Lemke CD, Salem AK. Nanoparticle delivery systems in cancer vaccines. *Pharm Res* 2011;**28**:215–36
89. Cohen EP, Chopra A, IOS, Kim TS. Enhancing cellular cancer vaccines. *Immunotherapy* 2009;**1**:495–504
90. Rosenberg SA, Yang JC, Restifo NP. Cancer immunotherapy: moving beyond current vaccines. *Nature Med* 2004;**10**:909–15
91. Altin JG, Parish CR. Liposomal vaccines – targeting the delivery of antigen. *Methods* 2006;**40**:39–52
92. Dubensky TW Jr., Reed SG. Adjuvants for cancer vaccines. *Semin Immunol* 2010;**22**:155–61
93. Lima KM, dos Santos SA, Rodrigues JM Jr., Silva CL. Vaccine adjuvant: it makes the difference. *Vaccine* 2004;**22**:2374–79
94. Winter H, Fox BA, Ruttinger D. Future of cancer vaccines. *Methods Mol Biol* 2014;**1139**:555–64
95. Fan Y, Moon JJ. Nanoparticle drug delivery systems designed to improve cancer vaccines and immunotherapy. *Vaccines (Basel)* 2015;**3**:662–85
96. Shao K, Singha S, Clemente-Casares X, Tsai S, Yang Y, Santamaria P. Nanoparticle-based immunotherapy for cancer. *ACS Nano* 2015;**9**:16–30
97. Wu TL, Ertl HC. Immune barriers to successful gene therapy. *Trends Mol Med* 2009;**15**:32–9
98. Irvine DJ, Swartz MA, Szeto GL. Engineering synthetic vaccines using cues from natural immunity. *Nat Mater* 2013;**12**:978–90
99. Gallucci S, Matzinger P. Danger signals: SOS to the immune system. *Curr Opin Immunol* 2001;**13**:114–19
100. Matzinger P. Tolerance, danger, and the extended family. *Annu Rev Immunol* 1994;**12**:991–1045
101. Bachmann MF, Rohrer UH, Kundig TM, Burki K, Hengartner H, Zinkernagel RM. The influence of antigen organization on B cell responsiveness. *Science* 1993;**262**:1448–51
102. Pasare C, Medzhitov R. Control of B-cell responses by toll-like receptors. *Nature* 2005;**438**:364–8
103. Shukla S, Wen AM, Commandeur U, Steinmetz NF. Presentation of HER2 epitopes using a filamentous plant virus-based vaccination platform. *J Mater Chem B* 2014;**2**:6249–58
104. Jobsri J, Allen A, Rajagopal D, Shipton M, Kanyuka K, Lomonosoff GP, Ottensmeier C, Diebold SS, Stevenson FK, Savelyeva N. Plant virus particles carrying tumour antigen activate TLR7 and induce high levels of protective antibody. *PLoS One* 2015;**10**:e0118096
105. Scheerlinck JP, Greenwood DL. Virus-sized vaccine delivery systems. *Drug Discov Today* 2008;**13**:882–7
106. Riedel T, Ghasparian A, Moehele K, Rusert P, Trkola A, Robinson JA. Synthetic virus-like particles and conformationally constrained peptidomimetics in vaccine design. *ChemBiochem* 2011;**12**:2829–36
107. Plummer EM, Manchester M. Viral nanoparticles and virus-like particles: platforms for contemporary vaccine design. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* 2010

108. Noad R, Roy P. Virus-like particles as immunogens. *Trends Microbiol* 2003;**11**:438–44
109. Fang RH, Kroll AV, Zhang L. Nanoparticle-based manipulation of antigen-presenting cells for cancer immunotherapy. *Small* 2015;**11**:5483–96
110. Joffre OP, Segura E, Savina A, Amigorena S. Cross-presentation by dendritic cells. *Nat Rev Immunol* 2012;**12**:557–69
111. Molino NM, Anderson AK, Nelson EL, Wang SW. Biomimetic protein nanoparticles facilitate enhanced dendritic cell activation and cross-presentation. *ACS Nano* 2013;**7**:9743–52
112. Saluja SS, Hanlon DJ, Sharp FA, Hong E, Khalil D, Robinson E, Tigelaar R, Fahmy TM, Edelson RL. Targeting human dendritic cells via DEC-205 using PLGA nanoparticles leads to enhanced cross-presentation of a melanoma-associated antigen. *Int J Nanomedicine* 2014;**9**:5231–46
113. Sharp FA, Ruane D, Claass B, Creagh E, Harris J, Malyala P, Singh M, O'Hagan DT, Petrilli V, Tschopp J, O'Neill LA, Lavelle EC. Uptake of particulate vaccine adjuvants by dendritic cells activates the NALP3 inflammasome. *Proc Natl Acad Sci U S A* 2009;**106**:870–5
114. Li P, Luo Z, Liu P, Gao N, Zhang Y, Pan H, Liu L, Wang C, Cai L, Ma Y. Bioreducible alginate-poly(ethylenimine) nanogels as an antigen-delivery system robustly enhance vaccine-elicited humoral and cellular immune responses. *J Control Release* 2013;**168**:271–9
115. Hu Y, Litwin T, Nagaraja AR, Kwong B, Katz J, Watson N, Irvine DJ. Cytosolic delivery of membrane-impermeable molecules in dendritic cells using pH-responsive core-shell nanoparticles. *Nano Lett* 2007;**7**:3056–64
116. Mukai Y, Yoshinaga T, Yoshikawa M, Matsuo K, Yoshikawa T, Matsuo K, Niki K, Yoshioka Y, Okada N, Nakagawa S. Induction of endoplasmic reticulum-endosome fusion for antigen cross-presentation induced by poly (gamma-glutamic acid) nanoparticles. *J Immunol* 2011;**187**:6249–55
117. Reddy ST, van der Vlies AJ, Simeoni E, Angeli V, Randolph CJ, O'Neil CP, Lee LK, Swartz MA, Hubbell JA. Exploiting lymphatic transport and complement activation in nanoparticle vaccines. *Nat Biotechnol* 2007;**25**:1159–64
118. Thomas SN, Vokali E, Lund AW, Hubbell JA, Swartz MA. Targeting the tumor-draining lymph node with adjuvanted nanoparticles reshapes the anti-tumor immune response. *Biomaterials* 2014;**35**:814–24
119. Rosalia RA, Cruz LJ, van Duikeren S, Tromp AT, Silva AL, Jiskoot W, de Gruijl T, Lowik C, Oostendorp J, van der Burg SH, Ossendorp F. CD40-targeted dendritic cell delivery of PLGA-nanoparticle vaccines induce potent anti-tumor responses. *Biomaterials* 2015;**40**:88–97
120. Hardy CL, Lemasurier JS, Mohamud R, Yao J, Xiang SD, Rolland JM, O'Hehir RE, Plebanski M. Differential uptake of nanoparticles and microparticles by pulmonary APC subsets induces discrete immunological imprints. *J Immunol* 2013;**191**:5278–90
121. Cruz LJ, Tacke PJ, Rueda F, Domingo JC, Albericio F, Figdor CG. Targeting nanoparticles to dendritic cells for immunotherapy. *Methods Enzymol* 2012;**509**:143–63
122. Krup OC, Kroll I, Bose G, Falkenberg FW. Cytokine depot formulations as adjuvants for tumor vaccines. I. Liposome-encapsulated IL-2 as a depot formulation. *J Immunother* 1999;**22**:525–38
123. Neelapu SS, Gause BL, Harvey L, Lee ST, Frye AR, Horton J, Robb RJ, Popescu MC, Kwak LW. A novel proteoliposomal vaccine induces antitumor immunity against follicular lymphoma. *Blood* 2007;**109**:5160–3
124. Hanes J, Sills A, Zhao Z, Suh KW, Tyler B, DiMeco F, Brat DJ, Choti MA, Leong KW, Pardoll DM, Brem H. Controlled local delivery of interleukin-2 by biodegradable polymers protects animals from experimental brain tumors and liver tumors. *Pharm Res* 2001;**18**:899–906
125. Sabel MS, Hill H, Jong YS, Mathiowitz E, Bankert RB, Egilmez NK. Neoadjuvant therapy with interleukin-12-loaded polylactic acid microspheres reduces local recurrence and distant metastases. *Surgery* 2001;**130**:470–8
126. Hill HC, Conway TF Jr, Sabel MS, Jong YS, Mathiowitz E, Bankert RB, Egilmez NK. Cancer immunotherapy with interleukin 12 and granulocyte-macrophage colony-stimulating factor-encapsulated microspheres: coinduction of innate and adaptive antitumor immunity and cure of disseminated disease. *Cancer Res* 2002;**62**:7254–63
127. Karkada M, Berinstein NL, Mansour M. Therapeutic vaccines and cancer: focus on DPX-0907. *Biologics* 2014;**8**:27–38
128. Eggermont LJ, Paulis LE, Tel J, Figdor CG. Towards efficient cancer immunotherapy: advances in developing artificial antigen-presenting cells. *Trends Biotechnol* 2014;**32**:456–65
129. van der Weijden J, Paulis LE, Verdoes M, van Hest JCM, Figdor CG. The right touch: design of artificial antigen-presenting cells to stimulate the immune system. *Chem Sci* 2014;**5**:3355–67
130. Sunshine JC, Perica K, Schneck JP, Green JJ. Particle shape dependence of CD8+ T cell activation by artificial antigen presenting cells. *Biomaterials* 2014;**35**:269–77
131. Zhang SX. Turning killer into cure – the story of oncolytic herpes simplex viruses. *Discov Med* 2015;**20**:303–9
132. Liu T, Li L, Fu C, Liu H, Chen D, Tang F. Pathological mechanisms of liver injury caused by continuous intraperitoneal injection of silica nanoparticles. *Biomaterials* 2012;**33**:2399–407
133. Chen Q, Xue Y, Sun J. Kupffer cell-mediated hepatic injury induced by silica nanoparticles in vitro and in vivo. *Int J Nanomed* 2013;**8**:1129–40
134. Moon C, Park HJ, Choi YH, Park EM, Castranova V, Kang JL. Pulmonary inflammation after intraperitoneal administration of ultrafine titanium dioxide (TiO₂) at rest or in lungs primed with lipopolysaccharide. *J Toxicol Environ Health A* 2010;**73**:396–409
135. Shvedova AA, Kisin ER, Mercer R, Murray AR, Johnson VJ, Potapovich AI, Tyurina YY, Gorelik O, Arepalli S, Schwegler-Berry D, Hubbs AE, Antonini J, Evans DE, Ku BK, Ramsey D, Maynard A, Kagan VE, Castranova V, Baron P. Unusual inflammatory and fibrogenic pulmonary responses to single-walled carbon nanotubes in mice. *Am J Physiol Lung Cell Mol Physiol* 2005;**289**:L698–708
136. Aggarwal P, Hall JB, McLeland CB, Dobrovolskaia MA, McNeil SE. Nanoparticle interaction with plasma proteins as it relates to particle biodistribution, biocompatibility and therapeutic efficacy. *Adv Drug Deliv Rev* 2009;**61**:428–37
137. Park K. Facing the truth about nanotechnology in drug delivery. *ACS Nano* 2013;**7**:7442–7
138. Blanco E, Shen H, Ferrari M. Principles of nanoparticle design for overcoming biological barriers to drug delivery. *Nat Biotechnol* 2015;**33**:941–51
139. Nel AE, Madler L, Velegol D, Xia T, Hoek EM, Somasundaran P, Klaessig F, Castranova V, Thompson M. Understanding biophysical-chemical interactions at the nano-bio interface. *Nat Mater* 2009;**8**:543–57
140. Pilla L, Rivoltini L, Patuzzo R, Marrari A, Valdagni R, Parmiani G. Multipeptide vaccination in cancer patients. *Expert Opin Biol Ther* 2009;**9**:1043–55
141. Morton SW, Herlihy KP, Shopsowitz KE, Deng ZJ, Chu KS, Bowerman CJ, Desimone JM, Hammond PT. Scalable manufacture of built-to-order nanomedicine: spray-assisted layer-by-layer functionalization of PRINT nanoparticles. *Adv Mater* 2013;**25**:4707–13
142. Steinmetz NF. Viral nanoparticles as platforms for next-generation therapeutics and imaging devices. *Nanomedicine* 2010;**6**:634–41
143. Pokorski JK, Steinmetz NF. The art of engineering viral nanoparticles. *Mol Pharm* 2011;**8**:29–43
144. DePorter SM, McNaughton BR. Engineered M13 bacteriophage nanocarriers for intracellular delivery of exogenous proteins to human prostate cancer cells. *Bioconjug Chem* 2014;**25**:1620–25
145. Shukla S, Eber FJ, Nagarajan AS, DiFranco NA, Schmidt N, Wen AM, Eiben S, Twyman RM, Wege C, Steinmetz NF. The impact of aspect ratio on the biodistribution and tumor homing of rigid soft-matter nanorods. *Adv Healthcare Mater* 2015;**4**:874–82
146. Cheng F, Tsvetkova IB, Khuong YL, Moore AW, Arnold RJ, Goicochea NL, Dragnea B, Mukhopadhyay S. The packaging of different cargo into enveloped viral nanoparticles. *Mol Pharm* 2013;**10**:51–58
147. Lee KL, Twyman RM, Fiering S, Steinmetz NF. Virus-based nanoparticles as platform technologies for modern vaccines. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* 2016; Epub ahead of print 19 January. DOI: 10.1002/wnan.1383